



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Inspection of immigrants at San Juan for the week ended January 11, 1902.

SAN JUAN, P. R., January 14, 1902.

SIR: I submit herewith report of alien steerage passengers arriving at this port during the week ended January 11, 1902. January 7, American schooner *India*, from Buen Ayre, Danish West Indies, with 1 immigrant.

Respectfully,

H. S. MATHEWSON,
Passed Assistant Surgeon, U. S. M. H. S.,
Chief Quarantine Officer for Porto Rico.

The SURGEON-GENERAL,
U. S. Marine-Hospital Service.

Foreign and insular statistical reports of countries and cities—Yearly and monthly.

AFRICA—*Sierra Leone*.—Week ended December 14, 1901. Estimated population, 35,000. Total number of deaths, 25. No contagious diseases reported.

BAHAMAS—*Dunmore Town*.—Two weeks ended January 17, 1902. Estimated population, 1,232. No deaths and no contagious diseases.

Governors Harbor.—Two weeks ended January 17, 1902. Estimated population, 1,575. No deaths and no contagious diseases.

Green Turtle Cay—Abaco.—Two weeks ended January 15, 1902. Estimated population, 3,314. No deaths and no contagious diseases.

Inagua.—Three weeks ended December 31, 1901. Estimated population, 1,460. Two deaths. No contagious diseases:

Nassau.—Two weeks ended January 20, 1902. Estimated population, 12,535. Number of deaths not reported. No contagious diseases reported.

CANADA—*British Columbia—Victoria*.—Month of December, 1901. Estimated population, 21,000. Total number of deaths, 20, including 2 from tuberculosis.

FRANCE—*Roubaix*.—Month of December, 1901. Estimated population, 124,660. Total number of deaths, 208, including enteric fever, 1, and 7 from whooping cough.

GERMANY—*Hanover*.—Month of November, 1901. Estimated population, 240,475. Total number of deaths, 241, including 8 from infectious diseases.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 75 great towns in England and Wales during the week ended January 4, 1902, correspond to an annual rate of 20.1 per 1,000 of the aggregate population, which is estimated at 14,800,427.

London.—One thousand eight hundred and ten deaths were registered during the week, including measles, 52; scarlet fever, 18; diphtheria, 24; whooping cough, 18; enteric fever, 7; smallpox, 28, and diarrhea and dysentery, 10. The deaths from all causes correspond to an annual rate of 20.6 per 1,000. In Greater London 2,482 deaths were registered, corresponding to an annual rate of 19.3 per 1,000 of the

population. In the "outer ring" the deaths included 12 from diphtheria, 10 from measles, 4 from scarlet fever, 10 from smallpox, and 5 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended January 4, 1902, in the 21 principal town districts of Ireland was 25.2 per 1,000 of the population, which is estimated at 1,092,322. The lowest rate was recorded in Lurgan, viz, 0.0, and the highest in Tralee, viz, 47.8 per 1,000. In Dublin and suburbs 223 deaths were registered, including diphtheria, 3; enteric fever, 2; whooping cough, 1, and 36 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended January 4, 1902, correspond to an annual rate of 23.2 per 1,000 of the population, which is estimated at 1,656,525. The lowest mortality was recorded in Leith, viz, 14.6, and the highest in Greenock, viz, 27.2 per 1,000. The aggregate number of deaths registered from all causes was 750, including diphtheria, 5; measles, 30; scarlet fever, 4, and 12 from whooping cough.

JAMAICA—Kingston.—Month of December, 1901. Estimated population, 46,542. Total number of deaths, 77, including 9 from phthisis pulmonalis.

Port Antonio.—Week ended January 11, 1901. Estimated population, not reported. Total number of deaths, not reported. The health of the port is good.

SPAIN—Barcelona.—Two weeks ended December 31, 1901. Estimated population, 600,000. Total number of deaths not reported. Thirteen deaths from enteric fever and 3 deaths from smallpox reported.

Corunna.—Month of December, 1901. Estimated population, 44,000. Total number of deaths, 140, including smallpox, 5, and 25 from tuberculosis.

Vigo.—Month of December, 1901. Estimated population, 20,000. Total number of deaths, 47, including 1 death from smallpox.

ST. HELENA.—Five weeks ended December 21, 1901. Estimated population, 3,342. Total number of deaths, 10, including 3 from tuberculosis.

WEST INDIES—St. Thomas.—Two weeks ended December 20, 1901. Estimated population, 11,012. Total number of deaths, 15. No contagious diseases.

Two weeks ended January 10, 1902. Total number of deaths, 14. No contagious diseases.